

(19) World Intellectual Property  
Organization  
International Bureau




(43) International Publication Date  
25 August 2005 (25.08.2005)

PCT

(10) International Publication Number  
**WO 2005/077005 A2**

- (51) International Patent Classification: **Not classified**
- (21) International Application Number:  
PCT/US2005/003877
- (22) International Filing Date: 4 February 2005 (04.02.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/542,000 4 February 2004 (04.02.2004) US
- (71) Applicant (for all designated States except US): **THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BLOCH, Kenneth, D.** [US/US]; 80 Park Street #32, Brookline, MA 02446 (US). **ICHINOSE, Fumito** [JP/US]; 60 Longwood Avenue, Apartment 803, Brookline, MA 02446 (US). **ZAPOL, Warren, M.** [US/US]; 30 Shepard St, Cambridge, MA 02138 (US). **EVGENOV, Oleg, V.** [NO/US]; 101 Rantoul Street, Apt.509, Beverly, MA 01915 (US).
- (74) Agent: **FRASER, Janis, K.**; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ENHANCING THE EFFECTIVENESS OF AN INHALED THERAPEUTIC GAS



(57) Abstract: Methods for enhancing the therapeutic or prophylactic effectiveness of an inhaled therapeutic gas are disclosed. The methods include administering to a mammal by inhalation a therapeutically effective amount of gaseous nitric oxide or carbon monoxide, and administering to the mammal a composition containing a compound that sensitizes soluble guanylate cyclase.

5

## ENHANCING THE EFFECTIVENESS OF AN INHALED THERAPEUTIC GAS

### Statement as to Federally Sponsored Research

This invention was made with Government support under grant numbers  
10 HL57172 and HL42397 awarded by the National Institutes of Health/National  
Heart, Lung, and Blood Institute. The Government may have certain rights in  
the invention.

### Cross Reference To Related Applications

15 This application claims priority from Provisional Application No.  
60/542,000, filed February 4, 2004. The prior application is incorporated herein  
by reference in its entirety.

### Field of the Invention

20 This invention relates to compositions and methods for enhancing the  
effectiveness of therapeutic gases.

### Background

Nitric oxide (NO) is a cell membrane-permeable, free radical molecule  
25 which accounts for the vasodilator activity of endothelium-derived relaxing  
factor (reviewed in Schmidt et al., Cell 78:919-925 [1994]). NO interacts with  
several intracellular molecular targets, one of which is soluble guanylate cyclase  
(sGC). Binding of NO to the heme group in sGC stimulates the conversion of  
guanosine triphosphate (GTP) to guanosine-3',5'-cyclic monophosphate (cGMP).  
30 cGMP exerts its effects on cells, in part, through its action on cGMP-dependent  
protein kinase (cGDPK). Additional cGMP targets include cGMP-gated ion  
channels and cGMP-regulated cyclic nucleotide phosphodiesterases.  
Phosphodiesterases (PDEs) inactivate cGMP by converting it to GMP.

The biological effects of NO are also mediated by cGMP-independent  
35 mechanisms. NO can serve as an antioxidant, opposing the effect of  
superoxides. The antioxidant properties of NO appear to account for its ability to  
modulate proinflammatory activation of endothelial cells. NO may also react

5 with superoxide to form peroxynitrite which may be responsible for the cellular toxicity associated with high levels of NO production.

### Summary

10 The invention is based, at least in part, on the discovery that a compound that sensitizes soluble guanylate cyclase can augment and/or prolong the therapeutic effectiveness of an inhaled therapeutic gas. The therapeutic gases described herein are nitric oxide and carbon monoxide.

In one aspect, the invention features a method for enhancing the therapeutic or prophylactic effectiveness of inhaled NO, the method including:  
15 (1) identifying a mammal (e.g., a human) that has or is at risk of developing a condition amenable to treatment or prevention by inhalation of gaseous NO; (2) administering to the mammal by inhalation a therapeutically effective amount of gaseous NO; and (3) administering to the mammal a composition containing a compound that sensitizes soluble guanylate cyclase, wherein the composition  
20 contains an amount of the compound sufficient to enhance the therapeutic or prophylactic effectiveness of the inhaled gaseous NO. In some embodiments, the method does not include the administration to the mammal of superoxide dismutase.

The invention also features a method for enhancing the therapeutic or  
25 prophylactic effectiveness of inhaled carbon monoxide (CO), the method including: (1) identifying a mammal (e.g., a human) that has or is at risk of developing a condition amenable to treatment or prevention by inhalation of gaseous CO; (2) administering to the mammal by inhalation a therapeutically effective amount of gaseous CO; and (3) administering to the mammal a  
30 composition containing a compound that sensitizes soluble guanylate cyclase, wherein the composition contains an amount of the compound sufficient to enhance the therapeutic or prophylactic effectiveness of the inhaled gaseous CO. In some embodiments, the method does not include the administration to the mammal of superoxide dismutase.

35 As detailed herein, inhalation of CO either alone or in combination with BAY 41-2272 (a compound that sensitizes soluble guanylate cyclase) was found to have no vasodilator effect on experimentally induced pulmonary

5 vasoconstriction. However, for those indications (e.g., treating or preventing  
ischemia-reperfusion injury or inflammation, extending the survival of an organ  
transplant, and inhibiting chronic rejection in a recipient) for which CO  
inhalation alone has been determined to have a beneficial effect, co-  
administration of a compound that sensitizes soluble guanylate cyclase is  
10 expected to enhance the therapeutic effectiveness of inhaled CO for those  
indications.

In some embodiments, prior to administering the therapeutic gas and the  
composition, the mammal is diagnosed as having pulmonary vasoconstriction.  
In other embodiments, prior to administering the therapeutic gas and the  
15 composition, the mammal is diagnosed as being at risk of developing pulmonary  
vasoconstriction. The pulmonary vasoconstriction can be, for example, acute  
pulmonary vasoconstriction, reversible pulmonary vasoconstriction, chronic  
pulmonary vasoconstriction which has a reversible component, or chronic  
pulmonary vasoconstriction which does not have a reversible component.

20 The mammal can have or be at risk of developing pneumonia, traumatic  
injury, aspiration or inhalation injury, fat embolism in the lung, acidosis,  
inflammation of the lung, adult respiratory distress syndrome, acute mountain  
sickness, post cardiac surgery acute pulmonary hypertension, persistent  
pulmonary hypertension of the newborn, perinatal aspiration syndrome, hyaline  
25 membrane disease, acute pulmonary thromboembolism, acute pulmonary edema,  
heparin-protamine reactions, sepsis, hypoxia, asthma, status asthmaticus, or  
hypoxia of the newborn.

The mammal can have or be at risk of developing chronic pulmonary  
hypertension, bronchopulmonary dysplasia, chronic pulmonary  
30 thromboembolism, idiopathic pulmonary hypertension, or chronic hypoxia.

In some embodiments, prior to administering the therapeutic gas and the  
composition, the mammal is diagnosed as having bronchoconstriction.

In other embodiments, prior to administering the therapeutic gas and the  
composition, the mammal is diagnosed as being at risk of developing  
35 bronchoconstriction. The bronchoconstriction can be, for example, associated  
with asthma.

5           In some embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as having a vascular thrombosis. In other embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as being at risk of developing a vascular thrombosis. The vascular thrombosis can be, for example, an arterial thrombosis or a venous thrombosis. In some cases, the vascular thrombosis is not a pulmonary thrombosis.

          In some embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as having or being at risk of developing an acute ischemic coronary syndrome. The acute ischemic coronary syndrome can be, for example, myocardial infarction, unstable angina pectoris, thrombosis after coronary revascularization, or reocclusion after coronary thrombolysis. The acute ischemic coronary syndrome can be associated with an artery-occluding disease. The acute ischemic coronary syndrome can be associated with a vascular interventional procedure (e.g., angioplasty such as percutaneous transluminal coronary angioplasty (PTCA), coronary artery bypass surgery, or a procedure to implant a coronary artery stent).

          In some embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as having arterial restenosis. In other embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as being at risk of developing arterial restenosis. In some cases, the mammal has undergone or is preparing to undergo a vascular interventional procedure (e.g., angioplasty such as PTCA, coronary artery surgery, or a procedure to implant a coronary artery stent).

          In those methods that involve a vascular interventional procedure to implant a stent, such a stent can optionally be coated with a compound such as an antiproliferative agent or a an agent that sensitizes soluble guanylate cyclase to NO (thereby sensitizing NO-exposed platelets and leukocytes which adhere to them).

          In some embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as having a hemoglobinopathy. In other embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as being at risk of developing a hemoglobinopathy.

5           The hemoglobinopathy can be characterized by (a) a reduced affinity of the patient's hemoglobin for oxygen compared with the affinity for oxygen of normal adult hemoglobin (Hb-A), or (b) a tendency of the patient's erythrocytes to sickle. In some cases, the hemoglobinopathy is selected from the group consisting of sickle cell trait; Hb-C, Hb-D, Hb-E, Hb-H, Hb-I, and Hb-Kansas  
10 disorders; or a combination of Hb-S with a second mutant  $\beta$ -globin allele. The hemoglobinopathy can be sickle cell disease.

          In some embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as having an ischemia-reperfusion injury. In other embodiments, prior to administering the therapeutic gas and the  
15 composition, the mammal is diagnosed as being at risk of developing ischemia-reperfusion injury. The ischemia-reperfusion injury can be in a non-pulmonary tissue. In some cases, the ischemia-reperfusion injury is caused by surgery, e.g., heart bypass surgery or transplantation surgery such as kidney transplantation surgery or heart transplantation surgery. In some cases, the ischemia-reperfusion  
20 injury occurs in the kidney, brain, or intestine. In some cases, the ischemia-reperfusion injury is caused by a stroke. In some cases, the ischemia-reperfusion injury occurs as a result of vascular occlusion at the time of aortic surgery. In other cases, the ischemia-reperfusion injury occurs as a result of re-vascularization of a limb.

25           In some embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as having inflammation. In other embodiments, prior to administering the therapeutic gas and the composition, the mammal is diagnosed as being at risk of developing inflammation. The inflammation can be in a non-pulmonary tissue. The inflammation can be  
30 associated with arthritis, myocarditis, encephalitis, transplant rejection, systemic lupus erythematosus, gout, dermatitis, inflammatory bowel disease, hepatitis, or thyroiditis.

          The invention also features a method of improving gas exchange in the lungs of a mammal, the method including: (1) identifying a mammal (e.g., a  
35 human) for whom an improvement in gas exchange within the lungs would be beneficial; (2) administering to the mammal by inhalation a therapeutically effective amount of gaseous NO; and (3) administering to the mammal a

5 composition containing a compound that sensitizes soluble guanylate cyclase, wherein the composition contains an amount of the compound sufficient to enhance the therapeutic effectiveness of the inhaled gaseous NO and improve gas exchange in the lungs of a mammal. In some embodiments, the method does not include the administration to the mammal of superoxide dismutase.

10 In some embodiments, the mammal is hypoxic and/or suffers from a lung injury.

The composition containing a compound that sensitizes soluble guanylate cyclase can be introduced into the mammal by, for example, an oral, intravenous, intramuscular, subcutaneous, or intraperitoneal route. In addition, 15 the composition can be introduced into the mammal by providing an aerosol or dry powder containing the composition for inhalation by the mammal. The composition can be inhaled in the therapeutic gas containing the gaseous NO or CO. Exemplary compounds that sensitize soluble guanylate cyclase are YC-1 (3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole) and BAY 41-2272 (3-(4-amino-5-cyclopropylpyrimidine-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4- 20 b]pyridine).

NO gas inhaled by the mammal can be administered at a predetermined concentration. Preferably it is administered in the absence of tobacco smoke. The predetermined concentration can be, for example, 0.1 ppm to 300 ppm, 1 25 ppm to 250 ppm, or 5 ppm to 200 ppm. Alternatively, the predetermined concentration can be, for example, at least 5 ppm, at least 40 ppm, at least 80 ppm, or 180 ppm or less.

In some embodiments, the therapeutic gas is inhaled continuously for an extended period or inhaled intermittently for an extended period. For example, 30 the therapeutic gas can be inhaled continuously for at least three minutes, after which inhalation can be stopped for a period of at least 0.5, 1, 2, 6, 12, or 24 hours prior to a subsequent inhalation.

An advantage of some of the methods described herein is the achievement of a desired therapeutic outcome by using an NO or CO dosage 35 lower than that required if NO or CO alone were administered to a patient. Lower dosages of NO or CO are expected to decrease the likelihood of adverse events that may accompany inhalation of higher doses of these therapeutic gases.

5           Another advantage of some of the methods described herein is that the co-administration of a compound that sensitizes soluble guanylate cyclase unexpectedly results in the prolongation of a response to an inhaled therapeutic gas. Accordingly, certain methods described herein can facilitate long-term NO or CO therapy by permitting an intermittent inhalation of gaseous NO or CO by  
10 a patient.

          Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Suitable methods and materials are described below, although methods and materials similar or equivalent to those  
15 described herein can also be used in the practice or testing of the present invention. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not  
20 intended to be limiting.

          Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

#### Brief Description of the Drawings

25           Figs. 1A-1B are graphs depicting the effects of BAY 41-2272 on mean pulmonary arterial pressure (1A) and pulmonary vascular resistance (1B).

          Figs. 2A-2B are graphs depicting the effects of L-NAME on pulmonary vasodilation (2A) and systemic vasodilation (2B) induced by BAY 41-2272.

          Figs. 3A-3D are graphs depicting percent changes of pulmonary arterial  
30 pressure (3A), percent changes of pulmonary vascular resistance (3B), half-time of reversal of pulmonary vasodilation ( $T_{1/2}$ ) (3C), and transpulmonary cGMP release during inhalation of NO alone (NO) or in combination with BAY 41-2272 (BAY + NO) (3D).

          Fig. 4 is a graph depicting the effects of YC-1, inhaled NO, and the  
35 combination of YC-1 and inhaled NO on pulmonary and systemic arterial pressure.



5                    Fig. 5 is a graph depicting the effects of inhaled NO and YC-1 administration on pulmonary vasodilation.

### Detailed Description

#### 10    Administration of Inhaled Nitric Oxide and Carbon Monoxide

                  Methods for safe and effective administration of NO and CO by inhalation are well known. See, e.g., Zapol, U.S. Patent No. 5,570,683; Zapol et al., U.S. Patent No. 5,904,938; Bach et al., U.S. Published Application No. 20030039638; and Frostell et al., 1991, *Circulation* 83:2038-2047. NO for  
15    inhalation is available commercially (INOmax™, INO Therapeutics, Inc., Clinton, NJ).

                  A suitable starting dosage for NO administered by inhalation may be 20 ppm. See, e.g., INOmax™ package insert. However, dosage can vary, e.g., from 0.1 ppm to 100 ppm, depending on the age and condition of the patient, the  
20    disease or disorder being treated, amount of sensitizer of soluble guanylate cyclase administered to the patient, and other factors that the treating physician may deem relevant. Preferably, the lowest effective dose is inhaled. To arrive at the lowest effective dosage empirically, administration can be commenced at 20 ppm and then decreased gradually until efficacy (e.g., vasodilator efficacy) is  
25    lost. Where 20 ppm is deemed an insufficient inhaled dose, NO dosage may be increased gradually until effectiveness (e.g., vasodilator effectiveness) is observed. Such adjustment of dosage is routine for those of skill in the art. An advantage of the present invention is that in many cases it enables achievement of a desired therapeutic outcome at an NO dosage lower than that required if NO  
30    were administered alone. In addition, it may allow for the prolongation of an inhaled NO response, which may allow for intermittent therapy (i.e., increased intervals between inhalations of NO by a subject).

                  Inhaled NO can be administered from a source of stored, compressed NO gas. The source of NO can be 100% NO, or diluted with N<sub>2</sub> or any other inert  
35    gas (e.g., helium). The NO can be obtained and stored as a mixture free of any contaminating O<sub>2</sub> or higher oxides of nitrogen, because such higher oxides of nitrogen (which can form by reaction of O<sub>2</sub> with NO) are potentially harmful to

5 lung tissues. If desired, purity of the NO may be demonstrated with chemiluminescence analysis, prior to administration to a patient. Chemiluminescence NO-NO<sub>x</sub> analyzers are commercially available (e.g., Model 14A, Thermo Environmental Instruments, Franklin, MA). The NO-N<sub>2</sub> mixture may be blended with air or O<sub>2</sub> through, for example, calibrated rotameters which  
10 have been validated previously with a spirometer. The final concentration of NO in the breathing mixture may be verified with a chemical or chemiluminescence technique (see, e.g., Fontijin et al., Anal. Chem. 42:575 (1970)). Alternatively, NO and NO<sub>2</sub> concentrations may be monitored by means of an electrochemical analyzer. Any impurities such as NO<sub>2</sub> can be scrubbed by exposure to NaOH  
15 solutions, baralyme, or sodalime. As an additional control, the FiO<sub>2</sub> of the final gas mixture may also be assessed. If desired, the ventilator may have a gas scavenger added to the expiratory outlet to ensure that significant amounts of NO will not escape into the adjacent environment.

In a hospital or emergency field situation, administration of NO gas  
20 could be accomplished, for example, by attaching a tank of compressed NO gas in N<sub>2</sub>, and a second tank of oxygen or an oxygen/N<sub>2</sub> mixture, to an inhaler designed to mix gas from two sources; by controlling the flow of gas from each source, the concentration of NO inhaled by the patient can be maintained at an optimal level. NO gas may also be mixed with room air, using a standard low-  
25 flow blender (e.g., Bird Blender, Palm Springs, CA). NO may be generated from N<sub>2</sub> and O<sub>2</sub> (i.e., air) by using an electric NO generator. Such a generator is described in Zapol U.S. Patent No. 5,396,882.

NO or CO may be provided intermittently from an inhaler. The use of an inhaler may be particularly advantageous if a compound that sensitizes soluble  
30 guanylate cyclase is administered, orally or by inhalation, in conjunction with the NO or CO.

Administration of a compound that sensitizes soluble guanylate cyclase may decrease the total dosage of NO or CO required (or allow intermittent dosage) to produce a satisfactory therapeutic or prophylactic effect. As a result  
35 of a decrease of NO or CO dosage or intermittent inhalations, an inhaler can be used less frequently. The compound that sensitizes soluble guanylate can be administered before (e.g., within 1, 12, or 24 hours before), during, or after (e.g.,

5        within 1, 12, or 24 hours after) inhalation of the gaseous NO or CO by the patient.

         Inhaled NO or CO can optionally be administered by nasal prongs, mask, tent, intra-tracheal catheter or endotracheal tube, for an extended period, i.e., days or weeks. The administration may be continuous, during the extended  
10       period. Alternatively, administration could be intermittent during the extended period. The administration of gaseous NO or CO may be via spontaneous or mechanical ventilation.

#### Compounds that Sensitize Soluble Guanylate Cyclase

15        NO decomposes rapidly by reacting with molecular oxygen to produce nitrite and nitrate. In addition, NO entering the blood is rapidly inactivated by tight binding to hemoglobin. For these reasons, NO has only a short half-life in arterial blood. As detailed herein, compounds that sensitize soluble guanylate cyclase (sGC) to NO activation can augment and prolong the action of inhaled  
20       NO.

         A compound that sensitizes soluble sGC can be introduced into a mammal by any suitable method, including via an oral, transmucosal, intravenous, intramuscular, subcutaneous, intraperitoneal, transcutaneous, or per rectum route. Alternatively, the compound can be inhaled by the mammal. For  
25       inhalation, the compound can be formulated as a dry powder or an aerosolized or nebulized solution having a particle or droplet size of less than 10  $\mu\text{m}$  for optimal deposition in the alveoli, and may optionally be inhaled in a therapeutic gas containing NO.

5           Compounds that sensitize soluble guanylate cyclase (sGC) to NO  
activation can be identified by routine assays. Both *in vitro* cell free and cell  
based assays can be used to determine if a compound, such as an sGC activator,  
can sensitize sGC to activation by NO. *In vitro* kinetic assays can be performed  
using purified sGC, e.g., sGC purified according to Humbert, et al., Eur. J.  
10   Biochem. 190:273–278 (1990). An *in vitro* assay can be used to determine  
whether a compound potentiates NO stimulation of sGC, e.g., by determining  
whether the compound causes a leftward shift of the NO concentration-response  
curve of sGC activity (decreases  $EC_{50}$ ) or increases the activity of sGC for any  
given dose of NO ( $V_{max}$ ). Examples of *in vitro* assays that can be used to show  
15   that a compound sensitizes sGC and causes a leftward shift of the concentration-  
response curve of sGC activity are described in Friebe et al., EMBO J. 15:6863–  
6868 (1996) and Friebe and Koesling, Mol. Pharmacol. 53:123–127 (1998).

Cell based assays employ cells with endogenous NO-sensitive sGC  
activity, such as platelets or aortic smooth muscle cells. In a platelet aggregation  
20   assay, the aggregation of stimulated platelets can be measured in the presence of  
an NO-donor alone, a candidate compound alone, or in the presence of both the  
candidate compound and the NO-donor. Platelet aggregation tracks cGMP  
production inside the cells, thus the synergistic effect of a compound and an NO-  
donor can be inferred from their combined effect on platelet aggregation. See,  
25   e.g., Friebe et al., Mol. Pharmacol. 54:962-967 (1998). Smooth muscle cell  
preparations can similarly be exposed to an NO-donor alone, to a compound  
alone, or to both the compound and the NO-donor. The accumulation of cGMP  
can then be measured to determine if a compound sensitizes sGC to an NO-  
donor. See, e.g., Mulsch et al., Brit. J. Pharmacol. 120:681-689 (1997).

5           Examples of compounds that sensitize sGC include 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole (YC-1; Russwurm, J. Biol. Chem. 277:24883-24888, (2002), Schmidt et al., Mol. Pharmac. 59:220-224 (2001), Friebe, et al., Mol. Pharmacol. 54:962-967, (1998)) and compounds loosely based on YC-1 such as the pyrazolopyridine BAY 41-2272 (Stasch et al., Nature 10 410:212-215 (2001), the BAY 41-2272 derivatives ortho-(BAY 50-6038), meta-(BAY 51-9491) and para-PAL-(BAY 50-8364) (Becker et al., BMC Pharmacol. 1:13 (2001)), and BAY 41-8543 (Brit. J. Pharmacol. 135:333-343 (2002)). Additional compounds that stimulate sGC activation include 1-benzyl-3-(substituted hetaryl)-fused pyrazole derivatives (U.S. Patent No. 6,180,656) and 15 heterocyclylmethyl-substituted pyrazole derivatives (U.S. Patent No. 6,166,027).

          Routine sGC activity assays can be carried out to determine if an sGC activator also functions to sensitize sGC to NO. Exemplary activators of sGC include the substituted isoindolone derivatives described in U.S. Patent No. 6,344,468, the sulfonylamino carboxylic acid N-arylamides described in U.S. 20 Patent No. 6,548,547, and the sulfur substituted sulfonylaminocarboxylic acid N-arylamides described in U.S. Patent No. 6,335,334.

#### Assessment of Effects of Inhaled Nitric Oxide

          When inhaled NO and a compound that sensitizes soluble guanylate cyclase are administered to treat or prevent a medical condition, it is in some 25 cases desirable to monitor the effects of the administrations. Such monitoring can be used, in a particular individual, to verify desirable effects of the treatment. Such monitoring is also useful in adjusting the dose level, duration, and frequency of administration of inhaled NO in a given individual.

          The effects of administration of inhaled NO and a compound that 30 sensitizes soluble guanylate cyclase on a patient can be assessed by standard medical analyses used to evaluate the condition to be treated. For example, if the patient suffers from pulmonary vasoconstriction, pulmonary artery pressure can be monitored via a flow-directed pulmonary artery catheter, cardiac ultrasound, or range-gated doppler techniques. In another example, if the patient 35 suffers from vascular thrombosis or arterial restenosis, these conditions can be monitored by examination of clinical manifestations such as chest pain,

5        electrocardiography, serial analyses of vascular patency by ultrasound, or  
coronary angiography.

#### Other Agents Administered with Inhaled Nitric Oxide

10        In some embodiments, a phosphodiesterase inhibitor can be administered  
in conjunction with NO inhalation to inhibit the breakdown of cGMP by  
endogenous phosphodiesterases (see, e.g., U.S. Patent Nos. 5,570,683 and  
5,823,180). The phosphodiesterase inhibitor can be introduced into the mammal  
by any suitable method, including via an oral, transmucosal, intravenous,  
intramuscular, subcutaneous or intraperitoneal route. Alternatively, the inhibitor  
15        can be inhaled by the mammal. A suitable phosphodiesterase inhibitor is  
Zaprinast (M&B 22948; 2-o-propoxyphenyl-8-azapurine-6-one; Rhone-Poulenc  
Rorer, Dagenham Essex, UK).

20        An antithrombotic agent can be administered together with NO in certain  
methods described herein (e.g., treatment or prevention of ischemia-reperfusion  
injury or vascular thrombosis). Such antithrombotic agents serve to restore  
perfusion of the tissues susceptible to ischemia-reperfusion injury via  
thrombolysis, and augment the therapeutic effects of inhaled NO by decreasing  
the potential for activation of platelets in non-pulmonary tissues. Examples of  
antithrombotic agents are aspirin, streptokinase, urokinase, tissue plasminogen  
25        activator ("t-PA"), met-t-PA (i.e., t-PA with an N-terminal methionine residue),  
FE1X (a t-PA analog), heparin, hirudin, Hirulog (a hirudin analog), ticlopidine,  
and IIb/IIIa (e.g., Rheopro). One or more such antithrombotic agents may be  
administered to a mammal before, during, or after treatment with inhaled NO, so  
that the potential of platelets to become activated in non-pulmonary tissues is  
30        decreased.

35        The following are examples of the practice of the invention. They are  
not to be construed as limiting the scope of the invention in any way.

#### Examples

5     Example 1: Pharmacological Sensitization of Soluble Guanylate Cyclase  
6     Produces Pulmonary Vasodilation and Modulates the Pulmonary Response to  
7     Inhaled Nitric Oxide In awake lambs instrumented with vascular catheters and a  
8     tracheostomy tube, the thromboxane analog U-46619 was infused intravenously  
9     to increase mean pulmonary arterial pressure (PAP) to 35 mm Hg. After a  
10    stabilization period, seven animals received an intravenous infusion of BAY 41-  
11    2272 (Alexis Biochemicals, Lausen, Switzerland) at 0.03, 0.1, and 0.3 mg·kg<sup>-1</sup>·h<sup>-1</sup>  
12    administered for 30 minutes each. In another eight animals, inhaled NO (2,  
13    10, and 20 ppm) was administered in random order for 10 minutes each followed  
14    by 15 minute NO-free periods. An intravenous infusion of BAY 41-2272 (0.1  
15    mg·kg<sup>-1</sup>·h<sup>-1</sup>) was then started. After increasing the infusion of U-46619 to  
16    maintain PAP at 35 mmHg, NO was inhaled as before BAY 41-2272. In an  
17    additional eight lambs, which received an intravenous infusion of the NO  
18    synthase inhibitor, L-NAME (25 mg/kg + 5 mg·kg<sup>-1</sup>·h<sup>-1</sup>), together with U-  
19    46619, NO was inhaled before and during the infusion of BAY 41-2272.

20           Data were analyzed using repeated measures ANOVA followed by  
21    Dunnett adjustment. A  $p < .05$  was considered statistically significant.

22           BAY 41-2272 produced dose-dependent reductions in mean PAP and  
23    pulmonary vascular resistance index (PVRI) that were significantly greater than  
24    the corresponding reductions in mean arterial pressure (MAP) and systemic  
25    vascular resistance index (SVRI) (Figs. 1A-1B). Data are mean ± SEM (n = 7)  
26    and represent percent changes from the pre-treatment pulmonary hypertension  
27    values. \*  $p < .05$  vs. MAP; †  $p < .05$  vs. SVRI.

28           L-NAME abolished systemic but not pulmonary vasodilation induced by  
29    BAY 41-2272 (0.1 mg·kg<sup>-1</sup>·h<sup>-1</sup>) (Figs. 2A-2B). Data are mean ± SEM (n = 8)  
30    and represent percent changes from the pre-treatment pulmonary hypertension  
31    values. \*  $p < .05$  vs. BAY; †  $p < .05$  vs. MAP; ‡  $p < .05$  vs. SVRI.

5           Inhaled NO produced a dose-dependent, selective pulmonary  
vasodilation (Figs. 3A-3B). After NO was discontinued, PAP rapidly returned to  
baseline pulmonary hypertension (PH) with a  $T_{1/2} < 1$  min (Fig. 3C). Inhaled  
NO at 20 ppm also increased  $PaO_2/FiO_2$  and reduced  $P(A-a)O_2$  and  $Qs/Qt$   
( $P < 0.05$ ). Inhalation of 10 and 20 ppm of NO increased transpulmonary cGMP  
10 release ( $P < 0.01$ ) (Fig. 3D), whereas there were no significant changes of arterial  
methemoglobin concentrations.

Administration of BAY 41-2272 markedly enhanced the reductions of  
PAP, PVRI, and PVRI/SVRI induced by inhaled NO ( $P < 0.05$ ) (Figs. 3A-3B).  
After NO was discontinued, the persistence of the pulmonary vasodilation (as  
15 reflected by  $T_{1/2}$ ) during BAY 41-2272 infusion was greater than that before the  
infusion ( $P < 0.05$ ) (Fig. 3C). In the presence of BAY 41-2272, NO inhaled at 10  
and 20 ppm produced minor reductions of pulmonary capillary wedge pressure  
(PCWP) and systemic vascular resistance index (SVRI) and increments in  
cardiac index (CI) and stroke volume index (SVI) ( $P < 0.05$  vs. baseline PH) and  
20 reduced right ventricle stroke work index (RVSWI) to a greater extent than did  
inhaled NO alone ( $P < 0.05$ ), whereas MAP, HR and central venous pressure  
(CVP) remained unchanged. Moreover, during BAY 41-2272 infusion,  
inhalation of 10 and 20 ppm NO augmented  $PaO_2/FiO_2$  and reduced  $P(A-a)O_2$   
and  $Qs/Qt$  ( $P < 0.01$  vs. baseline PH). The co-administration of BAY 41-2272  
25 and inhaled NO increased transpulmonary cGMP release to a greater extent than  
did inhaled NO alone ( $P < 0.05$ ) (Fig. 3D). During the period of NO  
administrations, the plasma BAY 41-2272 concentrations remained at a stable  
level.



5     Example 2: Sensitization of Soluble Guanylate Augments and Prolongs the  
      Effects of Inhaled Nitric Oxide

      A mouse model of pulmonary hypertension was used to examine the combined effects of YC-1 and inhaled NO. Three mice (average weight 25 g) were anesthetized, their chest opened, catheters were placed in the carotid and  
10    pulmonary arteries, and pulmonary and systemic arterial pressures were recorded (Fig. 4). U46619, a thromboxane analogue was administered intravenously to induce pulmonary vasoconstriction. NO (4 parts per million) was added to the inspired gas, and pulmonary vasodilation was measured. Thereafter, NO  
15    inhalation was discontinued, and the duration of the pulmonary vasodilator effect was measured.

      YC-1 was administered as boluses over one minute (10, 50 and 100 ug), and both pulmonary artery pressure and systemic blood pressure were measured. Thereafter, YC-1 was administered as an infusion (5 ug/min), and the rate of  
20    U46619 infusion was increased (2- to 3-fold) to re-establish pulmonary vasoconstriction. The pulmonary vasodilator response to breathing 4 ppm NO was measured, and duration of pulmonary vasodilation after discontinuing NO was assessed. Bolus administration of YC-1 induced both pulmonary and  
      systemic vasodilation (Fig. 4).

      Breathing NO after YC-1 administration induced a greater pulmonary  
25    vasodilator response than did breathing NO alone. Addition of NO inhalation to YC-1 did not cause systemic vasodilation. Moreover, the duration of pulmonary vasodilation after discontinuing NO inhalation was greater during an infusion of YC-1 than before ( $P<0.01$ ; Fig. 5).

5     Example 3: Effects of Carbon Monoxide Inhaled Alone or in Combination with  
      BAY 41-2272

      Inhalation of CO alone or following administration of BAY 41-2272 had  
no vasodilator effect on the U-46619-induced pulmonary vasoconstriction.  
Systemic hemodynamics, lung gas exchange, and transpulmonary cGMP release  
10    were also unchanged by CO inhalation. Arterial concentrations of  
carboxyhemoglobin gradually rose from  $1.0 \pm 0.2\%$  to  $4.7 \pm 0.4\%$  ( $P < 0.01$ ) and  
from  $1.4 \pm 0.1\%$  to  $5.2 \pm 0.2\%$  ( $P < 0.01$ ), respectively, after breathing 500 ppm CO  
alone or in combination with BAY 41-2272. The plasma levels of BAY 41-2272  
remained stable during the period of CO administrations.

15

Other Embodiments

      While the invention has been described in conjunction with the detailed  
description thereof, the foregoing description is intended to illustrate and not  
limit the scope of the invention, which is defined by the scope of the appended  
20    claims. Other aspects, advantages, and modifications are within the scope of the  
following claims.

      What is claimed is:

- 5           1. A method for enhancing the therapeutic or prophylactic effectiveness  
of inhaled nitric oxide, the method comprising:  
          identifying a mammal that has or is at risk of developing a condition  
          amenable to treatment or prevention by inhalation of gaseous nitric oxide;  
          administering to the mammal by inhalation a therapeutically effective  
10       amount of gaseous nitric oxide; and  
          administering to the mammal a composition comprising a compound that  
sensitizes soluble guanylate cyclase, wherein the composition comprises an  
amount of the compound sufficient to enhance the therapeutic or prophylactic  
effectiveness of the inhaled gaseous nitric oxide,  
15       wherein the method does not comprise the administration to the mammal  
of superoxide dismutase.
2. The method of claim 1, wherein, prior to administering the gaseous  
nitric oxide and the composition, the mammal is diagnosed as having or being at  
20       risk of developing pulmonary vasoconstriction.
3. The method of claim 1, wherein the mammal has or is at risk of  
developing pneumonia, traumatic injury, aspiration or inhalation injury, fat  
embolism in the lung, acidosis, inflammation of the lung, adult respiratory  
25       distress syndrome, acute mountain sickness, post cardiac surgery acute  
pulmonary hypertension, persistent pulmonary hypertension of the newborn,  
perinatal aspiration syndrome, hyaline membrane disease, acute pulmonary  
thromboembolism, acute pulmonary edema, heparin-protamine reactions, sepsis,  
hypoxia, asthma, status asthmaticus, or hypoxia of the newborn.  
30
4. The method of claim 1, wherein the mammal has or is at risk of  
developing chronic pulmonary hypertension, bronchopulmonary dysplasia,  
chronic pulmonary thromboembolism, idiopathic pulmonary hypertension, or  
chronic hypoxia.  
35

5           5. The method of claim 1, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing bronchoconstriction.

10           6. A method for enhancing the therapeutic or prophylactic effectiveness of inhaled nitric oxide, the method comprising:  
            identifying a mammal that has or is at risk of developing a non-pulmonary condition amenable to treatment or prevention by inhalation of gaseous nitric oxide;  
            administering to the mammal by inhalation a therapeutically effective  
15           amount of gaseous nitric oxide; and  
            administering to the mammal a composition comprising a compound that sensitizes soluble guanylate cyclase, wherein the composition comprises an amount of the compound sufficient to enhance the therapeutic or prophylactic effectiveness of the inhaled gaseous nitric oxide.

20           7. The method of claim 6, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing a vascular thrombosis.

25           8. The method of claim 6, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing an acute ischemic coronary syndrome.

30           9. The method of claim 6, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing arterial restenosis.

35           10. The method of claim 6, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing a hemoglobinopathy.

5           11. The method of claim 6, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing an ischemia-reperfusion injury.

10           12. The method of claim 6, wherein, prior to administering the gaseous nitric oxide and the composition, the mammal is diagnosed as having or being at risk of developing inflammation.

15           13. The method of any of claims 6-12, wherein the method does not comprise the administration to the mammal of superoxide dismutase.

15           14. A method of improving gas exchange in the lungs of a mammal, the method comprising:  
              identifying a mammal for whom an improvement in gas exchange within the lungs would be beneficial;  
20           administering to the mammal by inhalation a therapeutically effective amount of gaseous nitric oxide; and  
              administering to the mammal a composition comprising a compound that sensitizes soluble guanylate cyclase, wherein the composition comprises an amount of the compound sufficient to enhance the therapeutic effectiveness of  
25           the inhaled gaseous nitric oxide and improve gas exchange in the lungs of a mammal,  
              wherein the method does not comprise the co-administration of superoxide dismutase.

30           15. The method of claim 14, wherein the mammal is hypoxic.

              16. The method of any of claims 1-15, wherein the composition is inhaled in a gas comprising the gaseous nitric oxide.

35           17. A method for enhancing the therapeutic or prophylactic effectiveness of inhaled carbon monoxide, the method comprising:

5 identifying a mammal that has or is at risk of developing a condition  
amenable to treatment or prevention by inhalation of gaseous carbon monoxide;  
administering to the mammal by inhalation a therapeutically effective  
amount of gaseous carbon monoxide; and  
administering to the mammal a composition comprising a compound that  
10 sensitizes soluble guanylate cyclase, wherein the composition comprises an  
amount of the compound sufficient to enhance the therapeutic or prophylactic  
effectiveness of the inhaled gaseous carbon monoxide.

18. The method of claim 17, wherein, prior to administering the gaseous  
15 carbon monoxide and the composition, the mammal is diagnosed as having or  
being at risk of developing an ischemia-reperfusion injury.

19. The method of claim 17, wherein, prior to administering the gaseous  
carbon monoxide and the composition, the mammal is diagnosed as having or  
20 being at risk of developing inflammation.

20. The method of claim 17, wherein, prior to administering the gaseous  
carbon monoxide and the composition, the mammal is diagnosed as having a  
hemoglobinopathy.

25 21. The method of any of claims 17-20, wherein the method does not  
comprise the administration to the mammal of superoxide dismutase.

22. The method of any of claims 17-21, wherein the composition is  
30 inhaled in a gas comprising the gaseous carbon monoxide.

23. The method of any of claims 1-22, wherein the compound that  
sensitizes soluble guanylate cyclase is 3-(5'-hydroxymethyl-2'-furyl)-1-  
benzylindazole (YC-1).

35

- 5           24. The method of any of claims 1-22, wherein the compound that sensitizes soluble guanylate cyclase is 3-(4-amino-5-cyclopropylpyrimidine-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine (BAY 41-2272).
25. The method of any of claims 1-24, wherein the mammal is a human.

1/5




Fig. 1A




Fig. 1B



2/5




Fig. 2A




Fig. 2B

3/5




Fig. 3A




Fig. 3B




Fig. 3C




Fig. 3D

4/5




Fig. 4

5/5




Fig. 5

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 August 2005 (25.08.2005)

PCT

(10) International Publication Number  
**WO 2005/077005 A3**

(51) International Patent Classification:  
**A61K 31/04** (2006.01)

(21) International Application Number:  
PCT/US2005/003877

(22) International Filing Date: 4 February 2005 (04.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/542,000 4 February 2004 (04.02.2004) US

(71) Applicant (for all designated States except US): **THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BLOCH, Kenneth, D.** [US/US]; 80 Park Street #32, Brookline, MA 02446 (US). **ICHINOSE, Fumito** [JP/US]; 60 Longwood Avenue, Apartment 803, Brookline, MA 02446 (US). **ZAPOL, Warren, M.** [US/US]; 30 Shepard St, Cambridge, MA 02138 (US). **EVGENOV, Oleg, V.** [NO/US]; 101 Rantoul Street, Apt.509, Beverly, MA 01915 (US).

(74) Agent: **FRASER, Janis, K.**; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
31 May 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ENHANCING THE EFFECTIVENESS OF AN INHALED THERAPEUTIC GAS

(57) Abstract: Methods for enhancing the therapeutic or prophylactic effectiveness of an inhaled therapeutic gas are disclosed. The methods include administering to a mammal by inhalation a therapeutically effective amount of gaseous nitric oxide or carbon monoxide, and administering to the mammal a composition containing a compound that sensitizes soluble guanylate cyclase.



WO 2005/077005 A3

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US05/03877

## A. CLASSIFICATION OF SUBJECT MATTER

IPC: A61K 31/04( 2006.01)

USPC: 514/718

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/718

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,485,827 A (ZAPOL et al.) 23 January 1996 (23.01.1996), columns 1-5.	1-25
Y	US 5,904,938 A (ZAPOL et al.) 18 May 1999 (18.05.1999), columns 1-7.	1-25
Y	BECKER et al., BMC Pharmacology (2001), Vol. 1, No. 13, pages 1-12.	1-25



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T"
"A" document defining the general state of the art which is not considered to be of particular relevance	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 April 2007 (02.04.2007)

Date of mailing of the international search report

20 APR 2007

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (571) 273-3201

Authorized officer

Valerie Bell-Harris  
Frank Choi

Telephone No. (571) 272-1600

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US05/03877

Continuation of B. FIELDS SEARCHED Item 3:  
EAST  
search terms: nitric oxide, guanylate cyclase, superoxide dismutase